

Second Article: (Sharing with another inside and outside the specialization- Published).

Article title	Soil application of effective microorganisms and nitrogen alleviates salt stress in hot pepper (<i>Capsicum annuum</i> L.) plants.
Participants	<p>Abdelsattar Abdelkhalik¹, Taia A. Abd El-Mageed², Ibrahim A. A. Mohamed³, Wael M. Semida¹, Omar A. A. I. Al-Elwany¹, Ibrahim M. Ibrahim⁴, Khaulood A. Hemida⁵, Mohamed T. El-Saadony⁶, Synan F. AbuQamar^{7,*}, Khaled A. El-Tarabily^{7,8,9,*} and Mohammed A. H. Gyushi¹</p> <p>¹Horticulture Department, Faculty of Agriculture, Fayoum University, Fayoum, Egypt ²Soil and Water Department, Faculty of Agriculture, Fayoum University, Fayoum, Egypt ³Botany Department, Faculty of Agriculture, Fayoum University, Fayoum, Egypt ⁴Department of Agricultural Microbiology, Faculty of Agriculture, Fayoum University, Fayoum, Egypt ⁵Botany Department, Faculty of Science, Fayoum University, Fayoum, Egypt ⁶Department of Agricultural Microbiology, Faculty of Agriculture, Zagazig University, Zagazig, Egypt ⁷Department of Biology, College of Science, United Arab Emirates University, Al Ain, United Arab Emirates ⁸Khalifa Center for Genetic Engineering and Biotechnology, United Arab Emirates University, Al-Ain, United Arab Emirates ⁹Harry Butler Institute, Murdoch University, Murdoch, Western Australia, Australia</p>
Article status	Sharing with another inside and outside the specialization- Published in an international specialized journal
The Journal	Frontiers in Plant Science, 13:1079260, 1–18.
Impact Factor	4.1

Abstract

The applications of effective microorganisms (EMs) and/or nitrogen (N) have a stimulating role on plants against abiotic stress conditions. The aim of the present study was to determine the impact of the co-application of EMs and N on growth, physio-biochemical attributes, anatomical structures, nutrients acquisition, capsaicin, protein, and osmoprotectant contents, as well as the antioxidative defense system of hot pepper (*Capsicum annuum* L.) plants. In the field trials, EMs were not applied (EMs⁻) or applied (EMs⁺) along with three N rates of 120, 150, and 180 kg unit N ha⁻¹ (designated as N₁₂₀, N₁₅₀, and N₁₈₀, respectively) to pepper plants grown in saline soils (9.6 dS m⁻¹). The application of EMs and/or high N levels attenuated the salt-induced damages to pepper growth and yield. The application of EMs⁺ with either N₁₅₀ or N₁₈₀ increased the number, average weight and yield of fruits by 14.4 or 17.0%, 20.8 or 20.8% and 28.4 or 27.5%, respectively, compared to pepper plants treated with the recommended dose (EMs⁻ × N₁₅₀). When EMs⁺ was individually applied or combined with either N₁₅₀ or N₁₈₀, we observed increased accumulation of capsaicin by 16.7 or 20.8%, protein by 12.5 or 16.7%, proline by 19.0 or 14.3%, and total soluble sugars by 3.7 or 7.4%, respectively, in comparison with those treated with the integrative EMs⁻ × N₁₅₀. In addition, the non-enzymatic contents (ascorbate, and glutathione) and enzymatic activities (catalase, superoxide dismutase, and glutathione reductase) of the antioxidant defense systems significantly increased in pepper plants treated with EMs⁺ alone or combined with N₁₅₀ or N₁₈₀ under salt stress conditions. Higher accumulation of nutrients (N, P, K⁺, and Ca²⁺) along with reduced Na⁺ acquisition was also evidenced in response to EMs⁺ or/and high N levels. Most anatomical features of stems and leaves recovered in pepper plants grown in saline soils and supplied with EMs⁺ and N. The application of EMs

and N is undoubtedly opening new sustainable approaches toward enhancing *abiotic stress* tolerance in crops (e.g. hot pepper).