



جامعة الفيوم  
كلية التمريض

## UROLOGY

A) Choose the correct answer

1) Proximal ureteral and renal stones smaller than 2 cm is managed by.

- ESWL
- PCNL
- Ureteroscopy
- Laparscopy

2) The normal value of PSA is

- 10 ng/ml
- 0-4 ng/ml.
- 4-10 ng/ml
- zero

3) Metastatic prostatic disease is managed by

- Radical prostatectomy
- Chemotherapy
- Radiotherapy
- Hormonal therapy

4) Alpha blockers includes the following except

- Alfuzosin
- Finasteride
- Doxazosin
- Tamsulosin

5) The most accurate technique of localizing a nonpalpable testis is.

- Ultrasound
- CT
- Laparscopy
- MRI

B) Complete the following sentences:

1) Hypospadias is a common congenital anomaly of the penis in which the urethra opens.....

2) Patients with superficial bladder cancer treated by.....

3) Patients with localized prostate cancer treated by.....

4) Patients with metastatic renal cell carcinoma treated by.....

5) ..... is the gold standard for treatment of distal ureteral calculi

C) What are the indications of PCNL?

D) Give short account about cancer bladder staging

E) What are the indications of surgery in patients with BPH?

GOOD LUCK