



البحث السادس

عنوان البحث باللغة الانجليزية:

Comparison Between Ketamine and Dexmedetomidine in Achieving Opioid Sparing Sedoanalgesia in Patients Following Open Heart Surgery: A Randomized Clinical Trial

نوع البحث:

بحث مشترك منشور غير مشتق من رسالة علمية سبق تقييمه في لجنة ترقية د. عبير شعبان جودة لدرجة استاذ مساعد الدورة (14) بتاريخ 2025/2/28 وتم تقييمه مقبول و حصل على 63%.

المشاركون في البحث حسب الترتيب:

د. محمد أحمد حامد ، د. مصطفى محمد الحمامصي، د. أحمد محمد الدمرداش ، د. وليد سعد طلبية ، د. اسماعيل سيد الديب، د. مهدي احمد عبد الهادي ، د. عبير شعبان جودة، د. عمر سيد فرغلي ، د. رنا أحمد عبدالغفار، د. محمد احمد شوقي ، د. علياء عبد الستار ، د. مينا محروس صبحي ، د. محمد حسن رجب، د. محمد فؤاد الجيار و د. ياسر سالم مصطفى.

تاريخ و مكان النشر:

Anesthesia & Pain Research, 2024-07, ISSN: 2639-846X.

ملخص البحث السادس باللغة الانجليزية:

Background: Opioid-sparing sedoanalgesia relies on non-opioid multimodal analgesic and sedative agents. It is being developed to decrease or eliminate opioid use for the management of acute postoperative pain to avoid opioid complications. This study aimed to compare ketamine and dexmedetomidine for their efficiency and safety in inducing opioid-sparing sedoanalgesia in patients following open heart surgery.

Methods: Ninety adult cardiac patients who underwent open heart surgery were enrolled. They were randomly divided into three groups. Group D received a dexmedetomidine infusion of 0.1– 0.2 $\mu\text{g/kg/hour}$, group K received a ketamine infusion of 1–2 $\mu\text{g/kg/min}$ while group C received a placebo as a control group. Drugs were given after the operation in the surgical intensive care unit. Total postoperative opioid consumption was the primary outcome. Richmond agitation sedation scale, visual analog pain scale, time to first analgesic request, patient satisfaction, and incidence of any complications were recorded.

Results: In this study, we found that the median (interquartile range) postoperative opioid consumption was significantly lower in groups D and K than in group C (156 (40), 140 (83), 185 (110); $p < 0.001$). Richmond agitation sedation scale was statistically significant until 6 hours postoperative ($p\text{-value: } < 0.05$). Patient satisfaction score was higher in groups D and K than in group C ($p\text{-value: } 0.005$). On the other side, there is no statistically significant difference between the three study groups regarding time to first analgesic request



كلية الطب

قسم التخدير و الرعاية المركزة الجراحية و علاج الألم



جامعة الفيوم

(minutes) (p-value: 0.064) or visual analog pain scale (p-value: > 0.05). Incidence of complications was highly significant between the study groups (p-value: < 0.001).

Conclusions: To decrease postoperative opioid use with their multiple side effects and improve patient satisfaction scores in patients undergoing open heart surgery under general anesthesia, the use of either ketamine or dexmedetomidine is recommended in the postoperative period as an adjuvant modality and a good idea with observation of possibility of some preventable side effects.