

# Achieving Target IOP

## Abstract

**Purpose:** Evaluation of progression of glaucoma in patients after achieving their target IOP using SAIF target IOP table

**Subject and Methods:** This is a retrospective non randomized comparative interventional study that was performed on 216 eyes of 108 patients with Primary open angle (POAG).and Normal tension glaucoma from the outpatient clinics at Fayoum University hospital and MISR University hospital, from 2009 till 2014 . We calculated the target IOP according to Saif's table<sup>4</sup> of prediction and C/D ratio. Patients were classified into two groups:

**Group 1 :** achieved target IOP( 48 female& 24 male) patients.

**Group 2 :** didn't achieve target IOP (18 female& 18 male) patients.

The mean C/D ratio in group 1 was  $0.373 \pm 0.179$ , (ranged 0.3 - 0.8) , while group 2: the mean C/D ratio was  $0.860 \pm 0.103$  (ranged 0.6 -0.93)

The V.F. difference in group 1: the mean (MD) was  $-1.90 \pm 4.92$  , (ranged -16.60 to -1.90). and group 2: the mean (MD) was  $0.27 \pm 1.48$  (ranged . -1.90 to 0.03)

**Conclusion :**After comparing visual field (MD) difference between two groups we found that there is statistically significant difference between both groups as regard the group that achieved target IOP there was regressive changes or stabilization of the visual field MD . Optimal target IOP may be different for different individuals depending on the severity of the disease and should be updated periodically as the disease progress

DFB NM

**Keywords:** Target IOP, primary open angel glaucoma, normal tension glaucoma, glaucoma suspect

## Introduction

It is difficult to define glaucoma precisely, as it encompasses a diverse group of disorders. All forms of the disease have in common a potentially progressive and characteristic optic neuropathy which is associated with visual field loss as damage progresses, and in which intraocular pressure is usually a key modifying factor. Glaucoma is the second most prevalent eye condition, after cataract known to cause blindness worldwide<sup>(1)</sup>.

The actual etiology of the condition remains unknown <sup>(2)</sup> . Glaucoma consists	37
of many eye disorders, such as congenital glaucoma, secondary glaucoma, primary	38
angle closure glaucoma (PACG), normal tension glaucoma (NTG), pigmentary	39
glaucoma, and primary open-angle glaucoma (POAG). These disorders destroy the	40
optic nerve, leading to blindness <sup>(3)</sup> .	41
The risk factors for getting glaucoma include age, race, sex, heridity, family	42
history, systemic(Diabetes, Obesity, Hypertension, Hypotension, Arteriosclerosis and	43
Smoking) and socioeconomic factors as well as local factors (myopia, corneal	44
thickness and scleral rigidity) all will channel into disc damage for the systemic	45
factors and level of IOP for the local factors. So calculation of the combined	46
probability of getting glaucoma for these 2 factors alone will include all the above	47
mentioned variables <sup>(4-9)</sup> .	48
<b>Target IOP</b> can be defined as the intraocular pressure level which is	49
necessary to prevent glaucomatous damage of visual field and optic nerve head in an	50
individual patient, and hinder the progression of already established, structural or	51
functional deficits. The criteria to help choose the target IOP include; the morphology	52
of the optic nerve head, the performance and stability of the visual field, and the	53
overall physical health of the patient <sup>(10)</sup> .	54
The following are the main problems of Target IOP assessment:	55
1.It must be individualized to the patient and to each eye. No absolute level or	56
percentage change from baseline will be correct for the majority of our patients <sup>(11)</sup> .	57
2.It must be an accurate estimate <sup>(11)</sup> .	58
3.It needs to be determined in advance. However we can only confirm the	59
appropriateness of the chosen IOP level at a later date. Trial and error is an	60
unavoidable part of the process <sup>(11)</sup> .	61
It is generally assumed that aiming to achieve a Target IOP with at least a 30%	62
reduction from the initial pressure at which damage occurred is a useful starting point	63
<sup>(11)</sup> .	64
Determining the Target IOP:	65
The target intraocular pressure is a "best guess" level of IOP. Below which	66
further damage to the optic nerve is unlikely to occur. The estimate is based on the	67
initial level of IOP, degree of existing damage (optic nerve cupping, reserving power	68
of the optic nerve, visual field loss, nerve fiber layer thickness) age, presence of other	69
risk factors (diabetes and arteriosclerotic vascular diseases), rate of progression if	70
known, family history of glaucoma <sup>(7)</sup> .	71

In average patient, the European Glaucoma Society (EGS) recommends that an initial target intraocular pressure should be set at least 30% lower than the pressure at which the ocular damage originally occurred. The more advanced the glaucoma, the greater the number of risk factors and the greater the vascular components, the lower the target IOP should be. The target IOP also helps the physician to assess the success of the treatment. The earlier the target IOP reached the better the outcome for the patient. The target intraocular pressure should be reassessed periodically and lowered if progression, optic nerve hemorrhage, or increase in risk factors occurs<sup>(13,14)</sup>.

## **Aim of the study**

Evaluation of progression of glaucoma in patients after achieving their target IOP using SAIF target IOP table

### ***Patients and methods***

This is a retrospective non randomized comparative interventional study that was performed on two hundreds and sixteen eyes of one hundred and eight patients. The ethical committee approval done before seeing the patients' records

#### **PATIENT SELECTION:**

##### **Inclusion criteria:**

- Primary open angle (POAG).
- Normal tension glaucoma.

##### **Exclusion criteria:**

- Closed angle glaucoma patients.
- Secondary glaucoma patients.
- Any previous ocular surgery.

#### **Patients:**

All patients attending the outpatient clinics at Beni Suef University hospital Fayoum University hospital and MISR University hospital, from 2009 till 2014.

#### **Examinations:**

**All patients must had** Full ophthalmological examinations included,

- Visual acuity assessment by Snellen's Chart.

● IOP measurement by Goldman's applanation tonometer at least 8 visits	103
● Slit lamp examination and fundus examination for optic disc evaluation by 90D lens.	104 105
● 2 Visual field analysis was done (Humphrey& Octopus).	106
● OCT for evaluation of (C/D ratio).	107
● Full medical assessment.	108 109
<b><u>Treatment:</u></b>	110
In this study we depended on medical treatment. Patients were treated with the suitable line of treatment according to initial IOP of the patient and to maintain target IOP after reaching it.	111 112 113
Lines of treatment:	114
-Monotherapy: either	115
*Beta blocker (e.g: Timolol) or	116
*Alpha2 agonist (e.g: Brimonidine).	117
-Bitherapy:	118
* Beta blocker& Alpha2 agonist or	119
* Beta blocker& prostaglandin analogue (e.g:latanoprost) or	120
* Alpha2 agonist& prostaglandin analogue.	121
* Beta blocker& Carbonic anhydrase inhibitor.	122 123
-Triple therapy:	124
*Beta blocker& carbonic anhydrase inhibitor and	125
*Alpha2 agonist, or	126
*Prostaglandin analogue.	127
-Quadruple therapy:	128
*Beta blocker& carbonic anhydrase inhibitor and	129
*Alpha2 agonist and	130
*Prostaglandin analogue.	131
<b>Calculation of target IOP:</b>	132
We calculated the target IOP according to Saif's table <sup>4</sup> of prediction and C/D ratio.	133 134

According to calculation of target IOP of each eye patients were classified into two groups:

-Group(1): achieving target IOP.

-Group(2): not achieving target IOP; including patients who didn't

Table(1) shows SAIF target IOP guided by the C/D ratio<sup>(4)</sup>.

C/D ratio	0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
Target IOP mmHg	20	18	17	16	15	14	13	11	10	9	8

We analyzed mean deviation of the visual field to assess V.F. progression with treatment,

### Results

Data were statically described in terms of mean,  $\pm$  standard deviation SD, median and range, or frequencies (number of cases) and percentages when appropriate.

This study included one hundred and eight patients (two hundreds and sixteen eyes), (sixty six female patients& forty two male patients) with primary open angle (POAG) & normal tension glaucoma divided into:

**Group 1** : achieved target IOP( 48 female& 24 male) patients.

**Group 2** : didn't achieve target IOP (18 female& 18 male) patients.

This study was done on 108 patients (216 eyes): 61,1% female patient (72.7% achieved target IOP & 27.3% didn't achieve target IOP) while 38.9% male patient (57.1% achieved target IOP & 42.9% didn't achieve target IOP) .

Table(2), demographic data (sex)

	Female		males	
	Number(n)	(%)	Number(n)	(%)
Achieved IOP	48	72.7%	24	57.1%
Not-achieved IOP	18	27.3%	18	42.9%

N=number

The mean age among patients achieved target IOP was 39 years old  $\pm$ 15years, min. age was 20 years old, max. age was 62 years old& among patients didn't achieve target IOP mean was 50 years old  $\pm$ 12 years, min. age was 39years old, max. age was 72 years as shown in table 3

The mean visual acuity was  $0.632 \pm 0.310$  in patients who achieved target IOP, in patients who didn't achieve target IOP mean was  $0.435 \pm 0.292$ , with minimal visual acuity 0.05 & maximum visual acuity 1.00 as shown in table 3

The mean C/D ratio in group 1 was  $0.373 \pm 0.179$ , (ranged 0.3 - 0.8), while group 2: the mean C/D ratio was  $0.860 \pm 0.103$  (ranged 0.6 - 0.93) as shown in table 3 and figure 1.

Table(3) statistical analysis of the 2 groups.

group		Mean	Median	Standard Deviation	Minimum	Maximum
Achieved IOP	age	39	47	15	14	57
	Visual Acuity	.632	.700	.310	.050	1.000
	Cup disc ratio	.373	.300	.179	.100	.800
	Visual Field baseline	6.691	3.080	7.335	.600	26.370
	Visual Field last visit	4.794	2.265	6.515	0.000	26.200
	Visual field difference	-1.90	-1.03	4.92	-16.60	7.10
	IOP visit 1	16.12	17.00	3.47	11.00	29.00
	IOP visit 2	15.31	14.50	2.20	12.00	22.00
	IOP visit 3	14.04	14.00	2.32	9.00	21.00
	IOP visit 4	13.35	13.00	2.61	9.00	23.00
	IOP visit 5	14.23	14.00	1.72	10.00	17.00
	IOP visit 6	14.00	14.00	2.58	9.00	20.00
	IOP visit 7	12.38	12.00	2.03	9.00	16.00
	IOP visit8	11.38	12.00	1.97	8.00	16.00
	IOP difference	4.73	4.50	2.95	0.00	13.00
	IOP decrease %	27.71	28.99	13.62	0.00	52.94
	Target IOP	15.192	16.000	1.987	10.000	18.000
not achieved IOP	age	50	49	12	39	72
	Visual Acuity	.435	.400	.292	.050	1.000
	Cup disc ratio	.860	.900	.103	.600	1.000
	Visual Field baseline	17.687	19.875	9.981	1.200	30.370
	Visual Field last visit	17.957	19.020	9.335	3.500	29.080
	Visual field difference	.27	.03	1.48	-1.90	2.80
	IOP visit 1	19.80	20.00	7.27	8.00	31.00
	IOP visit 2	16.60	15.50	4.69	11.00	25.00
	IOP visit 3	13.80	12.50	3.09	11.00	22.00
	IOP visit 4	14.60	13.50	4.06	9.00	21.00
	IOP visit 5	16.80	17.00	4.41	10.00	24.00
	IOP visit 6	15.10	14.00	2.83	12.00	20.00
	IOP visit 7	13.70	12.00	3.88	9.00	20.00
	IOP visit8	14.70	14.00	2.26	10.00	18.00
	IOP difference	5.10	5.00	6.35	-6.00	13.00
	IOP decrease %	14.21	28.17	39.07	-75.00	46.15
	Target IOP	9.500	9.000	1.295	8.000	13.000

As regard IOP changes among 8 visits;

At the **1<sup>st</sup> visit** the mean IOP was  $16.12 \pm 3.47$  mmHg among the group that achieved target IOP and it was  $19.80 \pm 7.27$  mmHg among the group that didn't achieve target IOP.

Mean IOP at the **2<sup>nd</sup> visit** became  $15.31 \pm 2.20$  mmHg among the group of patients that achieved target IOP and it was  $16.60 \pm 4.69$  mmHg among the group that didn't achieve target IOP.

Mean IOP at the **3<sup>rd</sup> visit** became  $14.04 \pm 2.32$  mmHg among the group of patients that achieved target IOP and it was  $13.80 \pm 3.09$  mmHg among the group that didn't achieve target IOP.

Mean IOP at the **4<sup>th</sup> visit** became  $13.35 \pm 2.61$  mmHg among the group of patients that achieved target IOP and it was  $14.60 \pm 4.06$  mmHg among the group that didn't achieve target IOP.

Mean IOP at the **5<sup>th</sup> visit** became  $14.23 \pm 1.72$  mmHg among the group of patients that achieved target IOP and it was  $16.80 \pm 4.41$  mmHg among the group that didn't achieve target IOP.

Mean IOP at the **6<sup>th</sup> visit** became  $14.00 \pm 2.58$  mmHg among the group of patients that achieved target IOP and it was  $15.10 \pm 2.83$  mmHg among the group that didn't achieve target IOP.

Mean IOP at the **7<sup>th</sup> visit** became  $12.38 \pm 2.03$  mmHg among the group of patients that achieved target IOP and it was  $13.70 \pm 3.88$  mmHg among the group that didn't achieve target IOP.

Mean IOP at the **8<sup>th</sup> visit** (last visit) became  $11.38 \pm 1.97$  mmHg among the group of patients that achieved target IOP and it was  $14.70 \pm 2.26$  mmHg among the group that didn't achieve target IOP .

IOP changes shown in table 3& 4 and figure 2

As regard IOP difference between the initial visit & the 8<sup>th</sup> visit patients who achieved target IOP the mean was  $4.73 \pm 2.95$  with min. difference zero and max difference 13.00mm Hg, patients who didn't achieve target IOP the mean was  $5.10 \pm 6.35$  with min. difference 6.00 mmHg & max difference 13.00 mmHg, as shown table 4 and figure 3.

Table(4) IOP difference between the 8 visits.

			Paired Differences					t	df	Sig. (2-tailed)
			Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
						Lower	Upper			
Achieve d IOP	Pair 1	VF1 - VF2	1.896923	4.922700	.394131	1.118361	2.675485	4.813	155	.000
	Pair 2	IOP1 - IOP2	.80769	2.11939	.16969	.47249	1.14289	4.760	155	.000

	Pair 3	IOP2 - IOP3	1.26923	1.95876	.15683	.95944	1.57902	8.093	155	.000
	Pair 4	IOP3 - IOP4	.69231	1.81966	.14569	.40451	.98010	4.752	155	.000
	Pair 5	IOP4 - IOP5	-.88462	3.02668	.24233	-1.36331	-.40592	-3.650	155	.000
	Pair 6	IOP5 - IOP6	.23077	2.17862	.17443	-.11380	.57534	1.323	155	.188
	Pair 7	IOP6 - IOP7	1.61538	2.29575	.18381	1.25229	1.97847	8.788	155	.000
	Pair 8	IOP7 - IOP8	1.00000	2.32795	.18639	.63182	1.36818	5.365	155	.000
	Pair 9	IOP1 - IOP8	4.73077	2.95197	.23635	4.26389	5.19765	20.016	155	.000
not Achieved IOP	Pair 1	VF1 - VF2	-.270000	1.479872	.191051	-.652291	.112291	-1.413	59	.163
	Pair 2	IOP1 - IOP2	3.20000	7.39400	.95456	1.28993	5.11007	3.352	59	.001
	Pair 3	IOP2 - IOP3	2.80000	5.31324	.68594	1.42744	4.17256	4.082	59	.000
	Pair 4	IOP3 - IOP4	-.80000	3.57392	.46139	-1.72324	.12324	-1.734	59	.088
	Pair 5	IOP4 - IOP5	-2.20000	5.91121	.76313	-3.72703	-.67297	-2.883	59	.005
	Pair 6	IOP5 - IOP6	1.70000	2.81822	.36383	.97198	2.42802	4.673	59	.000
	Pair 7	IOP6 - IOP7	1.40000	3.80544	.49128	.41695	2.38305	2.850	59	.006
	Pair 8	IOP7 - IOP8	-1.00000	3.15691	.40756	-1.81552	-.18448	-2.454	59	.017
	Pair 9	IOP1 - IOP8	5.10000	6.35317	.82019	3.45880	6.74120	6.218	59	.000

207

209

As regard initial V.F patients achieved target IOP the mean was  $6.691 \pm 7.335$ , while patients didn't achieve target IOP the mean was  $17.687 \pm 9.981$  as shown in table 5.

210

211

212

213

214

Table(5), mean visual field of patients achieved target IOP& patients didn't achieve target IOP

215

216

	Target IOP	
	Achieved	Not achieved
MD of V.F.(1)	6.691	17.687
MD of V.F.(2)	4.794	17.957



P value	.000	.163
Mean difference	-1.90(±4.92)	0.27(±1.48)

As regard 2<sup>nd</sup> V.F at the 8<sup>th</sup> visit in patients achieved target IOP the mean was 4.794 ±6.515 and in patients didn't achieve target IOP the mean was 17.957 ±9.335 as shown as shown in figure 4.

As regard V.F. difference patients who achieved target IOP the mean was -1.90 ±4.92 with min. difference -16.60 and max. difference -1.90. As regard patients who didn't achieve target IOP the mean was 0.27 ± 1.48 with min. diff. -1.90 & max. diff. 0.03 as shown in figure 5.

As regard line of treatment there was four lines of treatment which were individualized according to each patient condition;

42 patients (38.8%) used monotherapy 39 patients(36.11%) of them achieve target IOP but 3 patients(2.8%) didn't achieve target IOP

36 patients (33.3%) used bitherapy 21patients(19.4%) of them achieve target IOP but 15patients(13.9%) didn't achieve target IOP

15 patients (13.89%) used triple therapy 12 patients(11.1%) of them achieved target IOP but 3 patients(2.8%) didn't achieve target IOP

15 patients (13.89%) used quadruple therapy 6 patients(5.6%)of them achieved target IOP but 9 patients(8.3%) didn't achieve target IOP

As shown in table 6

Table(6),percentage of patients achieved& didn't achieve target IOP with different lines of treatment

	Line of treatment							
	Mono-therapy		Bi-therapy		Triple-therapy		Quadruple-therapy	
	Count	%	Count	%	Count	%	count	%
Achieved	78	36.1%	42	19.4%	24	11.1%	12	5.6%
Not achieved	6	2.8%	30	13.9%	6	2.8%	18	8.3%

	243
	244
	245
A male patient 29	246
Years old	247
His right eye achieved the target IOP while the left eye didn't achieve the target IOP	248
as shown in figure 6-9	249
	250
<b>Discussion</b>	251
Glaucoma is a progressive serious disease that ends up with blindness. Early	252
detection and diagnosis are no more dilemma. The dilemma will be how to stop or	253
slow the progression of the disease <sup>(6)</sup> .	254
Assessment of glaucoma progression includes evaluation of three main items: IOP&	255
optic disc and visual field.	256
Target IOP is defined as the mean intraocular pressure obtained with treatment that	257
prevents further glaucomatous damage <sup>(15)</sup> .	258
The risk factors for getting glaucoma include age, sex, race, heredity, family	259
history, systemic (Diabetes, Obesity, Hypertension, Hypotension, Arteriosclerosis and	260
Smoking) and socioeconomic factors as well as local factors (Myopia, Corneal	261
thickness and Scleral rigidity) all will channel into disc damage for the systemic	262
factors and level of IOP for the local factors <sup>(4-9)</sup> .	263
For the above mentioned reasons we used in this study the Saif 's Table for the target	264
IOP.	265
The aim of this study was to evaluate the progression of glaucoma by the	266
visual field changes and Cup disc ratio after reaching the target IOP by glaucoma	267
medical treatment.	268
There were 108 patients included in this study with primary open angle	269
glaucoma which was less than other studies and clinical trials that included larger	270
number of patients as <b>Tanuja&amp; Rajiv</b> on 150 cases of POAG and Normotensive	271
glaucoma <sup>(16)</sup> , the <b>Collaborative Initial Glaucoma Treatment Study(CIGTS)</b> on	272
607 patients with newly detected simple glaucoma <sup>(17-21)</sup> , <b>Early Manifest Glaucoma</b>	273
<b>Trial (EMT)</b> on 225 patients with newly diagnosed open angle glaucoma <sup>(22,23)</sup> ,	274
<b>Advanced Glaucoma Intervention Study (AGIS)</b> on 591 patients with advanced	275
open angle glaucoma with poor medical control of IOP <sup>(24-27)</sup> , the <b>Collaborative</b>	276
<b>Normal Tension Glaucoma Study (CNTGS)</b> on 230 patients with normal tension	277

glaucoma <sup>(28,29)</sup> , and the <b>Ocular hypertension treatment study(OHTS)</b> on 1836	278
patients with ocular hypertension <sup>(30-35)</sup> .	279
The age of patients in this study ranged from 20 to 72 years which is relatively	280
similar to the <b>Collaborative Normal Tension Glaucoma Study (CNTGS)</b> as the age	281
of patients in that study ranged from 20-90 years <sup>(28,29)</sup> .	282
We had two groups of patients in this study the first group achieved our	283
calculated target IOP while the other group didn't achieve the target IOP in two or	284
more visits.	285
<b>Achieved target IOP group:</b>	286
The IOP for the achieved group ranged from 8-16 mmHg which was similar to	287
the <b>Advanced Glaucoma Intervention Study</b> as the target IOP was set at <18	288
mmHg and the patients with lower IOP were free from visual field impairment,	289
whereas those with higher values of IOP showed sustained visual field	290
deterioration <sup>(24-27)</sup> .	291
<b>Tanuja&amp; Rajiv</b> showed that cases with a follow up range of 14& less and 15-20	292
mmHg were stable <sup>(16)</sup> .	293
<b>Early Manifest Glaucoma Trial (EMT)</b> set target IOP using percent reduction and	294
concluded that 25% reduction from the initial pressure decreased risk of progression	295
by 25% <sup>(22,23)</sup> .	296
<b>Collaborative Normal Tension Glaucoma study(CNTGS)</b> said that patients with	297
normal tension glaucoma (IOP<20 mmHg) with IOP reduction 30% showed a 12%	298
rate of visual field impairment at 5 years <sup>(28,29)</sup> .	299
The mean C/D ratio was 0.37±0.179 (ranged 0.3-0.8) Which is slight larger	300
than the mean C/D ratio of normal population (0.26 ± 0.14 ranged from 0.0 to 0.7)	301
and less than the glaucomatous group (0.50 ± 0.23 ranged from 0.1 – 0.9) in Beni	302
Suef area <sup>(36-38)</sup> .	303
<b>Not achieved target IOP group:</b>	304
The IOP for this group ranged from 18-30 mmHg with deterioration of the	305
visual field which was similar to the <b>Advanced Glaucoma Intervention Study</b> as	306
the target IOP was set at <18 mmHg and the patients with higher values of IOP	307
showed sustained visual field deterioration <sup>(24-27)</sup> .	308
<b>Tanuja&amp; Rajiv</b> said that analysis of visual field and optic disc changes of cases with	309
a follow up range of >20 mmHg showed deterioration <sup>(16)</sup> .	310
<b>Early Manifest Glaucoma Trial (EMT)</b> Study patients were divided into two	311
groups. In one group, 25% reduction of intraocular pressure was attained treatment,	312
whereas the other group was left untreated. Glaucoma progression measured by visual	313

field impairment was statistically significantly greater in the group of untreated patients than in those with intraocular pressure reduction<sup>(22,39)</sup>.

**Collaborative Normal Tension Glaucoma study(CNTGS)** said that patients with normal tension glaucoma (IOP<20 mmHg) left without treatment showed a 35% rate of progression of glaucomatous visual field impairment at 5 years<sup>(28,29)</sup>.

The visual field changes showed decrease in the MD as the mean of MD was 6.691 before treatment and became 4.794 after achieving target IOP among the group that achieved target IOP. This may be due to removal of the pressure from the ganglion cells and optic nerve, also short duration between the visual fields (6months to 3.5 years) between the study groups may be a factor in these visual field improvements.

Musch DC et al<sup>(20)</sup> showed a, substantial visual field loss and improvement over 5 years of follow-up In the collaborative initial glaucoma treatment study.

In the non-achieving group the MD was 17.687and became 17.957 even with treatment in the group not achieving target IOP, this was not shown in other studies that demonstrating variable changes and progression of the visual field<sup>(24,26,29,32,33,35)</sup>

## Conclusion

After comparing visual field (MD) difference between two groups we found that there is statistically significant difference between both groups as regard the group that achieved target IOP there was regressive changes or stabilization of the visual field MD

Optimal target IOP may be different for different individuals depending on the severity of the disease and should be updated periodically as the disease progress

The information gained from the study, assist in estimating and modifying target IOP.

## References

- 1- **Tamura H, Kawakami H, Kanamoto T.** High frequency of open-angle glaucoma in Japanese patients with Alzheimer's disease, J Neurol Sci,246:79–83,(2006)
- 2- **Dipiro J, Talbert RL, Yee GC, Wells BG, Posey LM.** Pharmacotherapy: A Pathophysiologic Approach,7th ed. New York, NY: McGraw Hill, (2009).
- 3- **Heiting G, Haddrill M.** Glaucoma: Types, Symptoms, Diagnosis and Treatment Available from <http://www.allaboutvision.com/conditions/glaucoma.htm# types of glaucoma>. Accessed June 6,(2015).

<b>4- Saif SSEH, Saif MYS, Saif AT.</b> Early Detection of Glaucoma, A New Scoring System. Bull Ophthalmol Soc Egypt, 2005; Vol 98, Number 3, 351-358.	348 349
<b>5- Saif SSEH, Saif MYS, Saif AT;</b> Target IOP What is New; Bull Ophthalmol Soc Egypt, 2006; Vol 99, Number 3, 445-449	350 351
<b>6- Saif SSEH, Saif MYS, Saif AT;</b> Glaucoma is it still a dilemma in the 21 <sup>st</sup> century; Bull Ophthalmol Soc Egypt, 2007; Vol 100, Number 3, 395-399	352 353
<b>7- Saif SSEH, Saif MYS, Saif AT;</b> Early Detection and Managment of Glaucoma , A New Scoring System. Highlights of Ophthalmology 2007; Volume 35, Number 6, 2-4	354 355 356
<b>8- Saif SSEH, Saif MYS, Saif AT;</b> The Glaucoma suspect, the dilemma. What is new? Bull Ophthalmol Soc Egypt, 2008; 101,	357 358
<b>9- Saif MYS, Saif AT, Saif PS, Salah ElDen W,</b> The Glaucoma suspect , the dilemma. What is new? Research in Ophthalmology 2013; 2(1): 10-14 doi:10.5923/j.ophtal	359 360 361
<b>10- Burk R, Volcker HE.</b> The concept of the" target IOP" in POAG. What level of intraocular pressure are we aiming at for which type of optic nerve head, Glaucoma World No.5(June1997).	362 363 364
<b>11- Traverso CE.</b> Practical points on the target IOP, Glaucoma World No.19(May2000).	365 366
<b>12- Nassar MK.</b> Threshold Amsler Grid testing to detect reserving power of the optic nerve,Bull. Ophthalmol. Soc. Egypt, 90:97-101(1997).	367 368
<b>13- EGS Terminology and Guidelines for Glaucoma 4th edition, 2014; 3.2, 134-138</b>	369
<b>14- Azarbod, P., Crawley, L., Ahmed, F., Cordeiro, M. F., &amp; Bloom, P.</b> (2015). Recent advances in the diagnosis and management of glaucoma. <i>Prescriber</i> ,26(1-2), 21-25.	370 371
<b>15-Nassar MK.</b> Target Intraocular Pressure. Available from <a href="http://www.glaucoma-egypt.org/Glaucoma-news/200102.pdf">http://www.glaucoma-egypt.org/Glaucoma-news/200102.pdf</a> . accessed (sep 2015).	372 373
<b>16-Kate T, Choudhary R.</b> Validity of Target IOP, AIOC proceedings, Glaucoma session II 231-232,(2005).	374 375
<b>17-Lichter PR, Musch DC, Gillespie BW.</b> Interim clinical outcomes in the Collaborative Initial Glaucoma Treatment Study comparing initial treatment randomized to medications or surgery. <i>Ophthalmology</i> ,108:1943-1953, 56,(2001).	376 377 378 379
<b>18- Musch, D. C., Gillespie, B. W., Niziol, L. M., Lichter, P. R., Varma, R., &amp; CIGTS Study Group.</b> (2011). Intraocular pressure control and long-term visual field loss in the Collaborative Initial Glaucoma Treatment Study. <i>Ophthalmology</i> , 118(9), 1766-1773.	380 381 382
<b>19- Musch, D., Niziol, L., &amp; Gillespie, B.</b> (2013). Factors associated with the onset of treatment in fellow eyes in the Collaborative Initial Glaucoma Treatment Study. <i>Investigative Ophthalmology &amp; Visual Science</i> , 54(15), 3484-3484.	383 384 385

<b>20-</b> Musch, D. C., Gillespie, B. W., Palmberg, P. F., Spaeth, G., Niziol, L. M., & Lichter, P. R. (2014). Visual Field Improvement in the Collaborative Initial Glaucoma Treatment Study. <i>American journal of ophthalmology</i> , 158(1), 96-104.	386 387 388
<b>21-</b> Musch, D. C., Niziol, L. M., Gillespie, B. W., & Lichter, P. R. (2015). An evaluation of target intraocular pressure use in the Collaborative Initial Glaucoma Treatment Study. <i>Investigative Ophthalmology &amp; Visual Science</i> , 56(7), 3693-3693.	389 390 391
<b>22-Leske MC, Heijl A, Hussein M.</b> Factors for glaucoma progression and the effect of treatment: the early manifest glaucoma trial. <i>Arch Ophthalmol.</i> , 121:48-56, (2003).	392 393 394
<b>23</b> Velpandian, T., Kotnala, A., Halder, N., Ravi, A. K., Archunan, V., & Sihota, R. (2014). Stability of Latanoprost in Generic Formulations Using Controlled Degradation and Patient Usage Simulation Studies. <i>Current eye research</i> , 40(6), 561-571.	395 396 397
<b>24-Kass MA, Gordon MO.</b> Intraocular pressure and visual field progression in open angle glaucoma, <i>Am. J. Ophthalmol.</i> , 130(4):4901, (2000, Oct.).	398 399
<b>25-</b> Zhang, L., Musch, D. C., Niziol, L. M., & Stein, J. D. (2012). Impact of Trabeculectomy Surgery on Global Visual Field Indices Using Data from Advanced Glaucoma Intervention Study (AGIS). <i>Investigative Ophthalmology &amp; Visual Science</i> , 53(14), 5950-5950.	400 401 402 403
<b>26-</b> Gros-Otero, J., Castejón, M., Paz-Moreno, J., Mikropoulos, D., & Teus, M. (2014). Perimetric progression using the Visual Field Index and the Advanced Glaucoma Intervention Study score and its clinical correlations. <i>Journal of optometry</i> .	404 405 406
<b>27-</b> Patel, P., Pritz, N., Fechtner, R., & Khouri, A. (2013). Assessment of Intervisit IOP Control in Eyes with Advanced Glaucoma. <i>Investigative Ophthalmology &amp; Visual Science</i> , 54(15), 5634-5634.	407 408 409
<b>28-Drance S, Andreson DR.</b> Collaborative Normal Tension Glaucoma Study Group, <i>Am. J. Ophthalmol.</i> , (2001).	410 411
<b>29-</b> Kim, M., Kim, D. M., Park, K. H., Kim, T. W., Jeoung, J. W., & Kim, S. H. (2013). Intraocular pressure reduction with topical medications and progression of normal-tension glaucoma: a 12-year mean follow-up study. <i>Acta ophthalmologica</i> , 91(4), e270-e275.	412 413 414 415
<b>30-Palmberg, P.,</b> Ocular Hypertension Treatment Study, <i>Arch. Ophthalmol.</i> (2002).	416
<b>31-</b> Mansberger, S. L., Gordon, M. O., Jampel, H., Bhorade, A., Brandt, J. D., Wilson, B., ... & Ocular Hypertension Treatment Study Group. (2012). Reduction in intraocular pressure after cataract extraction: the Ocular Hypertension Treatment Study. <i>Ophthalmology</i> , 119(9), 1826-1831.	417 418 419 420
<b>32-</b> De Moraes, C. G., Demirel, S., Gardiner, S. K., Liebmann, J. M., Cioffi, G. A., Ritch, R., ... & Ocular Hypertension Treatment Study Group. (2012). Rate of visual field progression in eyes with optic disc hemorrhages in the ocular hypertension treatment study. <i>Archives of ophthalmology</i> , 130(12), 1541-1546.	421 422 423 424
<b>33-</b> De Moraes, C. G., Demirel, S., Gardiner, S. K., Liebmann, J. M., Cioffi, G. A., Ritch, R., ... & Kass, M. A. (2012). Effect of treatment on the rate of visual field change in the ocular hypertension treatment study observation group. <i>Invest Ophthalmol Vis Sci</i> , 53(4), 1704-1709.	425 426 427 428

<b>34-</b> Brandt, J. D., Gordon, M. O., Gao, F., Beiser, J. A., Miller, J. P., Kass, M. A., & Ocular Hypertension Treatment Study Group. (2012). Adjusting intraocular pressure for central corneal thickness does not improve prediction models for primary open-angle glaucoma. <i>Ophthalmology</i> , 119(3), 437-442.	429 430 431 432
<b>35-</b> Demirel, S., De Moraes, C. G. V., Gardiner, S. K., Liebmann, J. M., Cioffi, G. A., Ritch, R., ... & Kass, M. A. (2012). The rate of visual field change in the ocular hypertension treatment study. <i>Investigative ophthalmology &amp; visual science</i> , 53(1), 224.	433 434 435
<b>36- Saif MYS , Khalil HMA, Abd El-Khalek MO, Makar A ;</b> Variations of Cup-to-Disc ratio in Age Group (18-40) Years Old . Research in Ophthalmology 2013; 2(1): 4-9 doi:10.5923/j.ophtal.(2013)0201.02	436 437 438
<b>37- Amer SAK, Saif MYS, Saif ATS, Saif PS;</b> Variations of Cup-to-Disc ratio in Children. The Open Ophthalmology Journal; 4:12-17,(2014).	439 440
<b>38- Saif MYS, Saif SEH, Saif ATS, Saif PS , Saftawy HS , Dabbous OA, Nabil M</b> . Evaluation of Large Cupping in Children with Anaemia Egyptian Ophthalmology Society dec 2015.EPUB	441 442 443
<b>39-</b> Gerometta, R., Escobar, D., & Candia, O. A. (2011). An hypothesis on pressure transmission from anterior chamber to optic nerve. <i>Medical hypotheses</i> , 77(5), 827-831.	444 445 446